



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\text{Then } EF = \frac{2(a^2 + ab - 2ab\sin^2\gamma)}{(a+b)\sqrt{1-e^2\sin^2\gamma}}.$$

$$\therefore EF = \frac{2a}{\sqrt{1-e^2\sin^2\gamma}} + \frac{2}{e\sqrt{ab}}\sqrt{1-e^2\sin^2\gamma} - \frac{2}{e\sqrt{ab}\sqrt{1-e^2\sin^2\gamma}}.$$

$$\therefore \int EF d\gamma = \frac{2}{e\sqrt{ab}} \{ (ae\sqrt{ab} - 1)F(e, \gamma) + E(e, \gamma) \}.$$

$$\therefore A = \frac{2}{\beta e\sqrt{ab}} \{ (ae\sqrt{ab} - 1)F_0^\beta(e, \gamma) + E_0^\beta(e, \gamma) \}, \quad b > a.$$

$$A = \frac{2}{\delta e\sqrt{ab}} \{ (ae\sqrt{ab} - 1)F_0^\delta(e, \gamma) + E_0^\delta(e, \gamma) \}, \quad b < a.$$

$$\text{II. } A_1 = 2 \int_0^{\theta'} \sqrt{a^2 - b^2 \sin^2 \theta} d\theta / \int_0^{\theta'} d\theta = \frac{2}{\theta'} \int_0^{\theta'} \sqrt{a^2 - b^2 \sin^2 \theta} d\theta.$$

$$\text{Let } \theta = \frac{1}{2}\pi + \lambda. \quad \therefore \theta' - \frac{1}{2}\pi = \lambda \text{ to } -\frac{1}{2}\pi = \lambda.$$

$$\begin{aligned} \therefore A_1 &= \frac{2}{\theta'} \int_{-\frac{1}{2}\pi}^{\theta' - \frac{1}{2}\pi} \sqrt{a^2 - b^2 \cos^2 \lambda} d\lambda = \frac{2}{\theta'} \int_{-\frac{1}{2}\pi}^{\theta' - \frac{1}{2}\pi} \sqrt{b^2 \sin^2 \lambda - (b^2 - a^2)} d\lambda \\ &= \frac{2\sqrt{(b^2 - a^2)}}{\theta'} \int_{-\frac{1}{2}\pi}^{\theta' - \frac{1}{2}\pi} \sqrt{\frac{b^2}{b^2 - a^2} \sin^2 \lambda - 1} d\lambda \\ &= \frac{2\sqrt{(b^2 - a^2)}}{\theta'} H_{-\frac{1}{2}\pi}^{\theta' - \frac{1}{2}\pi} \left(\frac{b}{\sqrt{b^2 - a^2}}, \lambda \right), \quad b > a. \end{aligned}$$

$$A_1 = 2 \int_0^{\frac{1}{2}\pi} \sqrt{a^2 - b^2 \sin^2 \theta} d\theta / \int_0^{\frac{1}{2}\pi} db = \frac{4a}{\pi} E_{\frac{1}{2}\pi}^{\frac{1}{2}\pi} \left(\frac{b}{a}, \theta \right), \quad b < a.$$

45. Proposed by J. C. WILLIAMS, Boston, Massachusetts.

At the end of the fifth inning the base ball score stands 7 to 9. What is the probability of winning for either team?

Solution by J. SCHEFFER, A. M., Hagerstown, Maryland.

From the stated score we are able to estimate the respective skill of the two teams, and their respective probabilities of winning the game.

The respective probabilities are $\frac{7}{18}$ and $\frac{9}{18}$. We have now to find the probabilities of either team winning at least 3 games out of 4, granting, of course, 9

innings to be played. These probabilities are respectfully, $(\frac{1}{16})^4 + 4(\frac{1}{16})^3 \cdot \frac{9}{16} = \frac{14}{16^4}$, and $(\frac{9}{16})^4 + 4(\frac{9}{16})^3 \cdot \frac{1}{16} = \frac{26}{16^4}$.

46. Proposed by J. A. CALDERHEAD, M. Sc., Professor of Mathematics in Curry University, Pittsburg, Pennsylvania.

Four men starting from random points on the circumference of a circular field and traveling at different rates, take random straight courses across it; find the chance that at least two of them will meet.

Professor Heaton says: "If the men are considered points the chance is 0." [A possible though difficult problem could be made of this one by using instead of men segments of straight lines moving along random secants of a circle, the velocity of the segments all being different. Editor.]

47. Proposed by HENRY HEATON, M. Sc., Atlantic, Iowa.

What is the average length of the chords that may be drawn from one extremity of the major axis of an ellipse to every point of the curve?

Solution by the PROPOSER.

The length of a single chord is

$$[(a-x)^2 + y^2]^{\frac{1}{2}} = (1/a)[a^2(a^2 - x^2) + b^2(a^2 - x^2)]^{\frac{1}{2}}.$$

Put S = distance around the ellipse. Then the required average is $A =$

$$\begin{aligned} \frac{2}{aS} \int_0^{180} [a^2(a-x)^2 + b^2(a^2 - x^2)]^{\frac{1}{2}} dS = \\ \frac{2}{a^2S} \int_{-a}^{+a} \frac{[a^2(a-x)^2 + b^2(a^2 - x^2)]^{\frac{1}{2}} [a^2(a^2 - x^2) + b^2x^2]^{\frac{1}{2}} dx}{(a^2 - x^2)^{\frac{1}{2}}} = \\ \frac{2}{a^2S} \int_{-a}^{+a} \frac{[a(a^2 + b^2) - (a^2 - b^2)x]^{\frac{1}{2}} [a^4 - (a^2 - b^2)x^2]^{\frac{1}{2}} dx}{(a+x)^{\frac{1}{2}}} \end{aligned}$$

This is readily reducible to elliptic functions of the first and second order, but the expressions I have been able to obtain are involved radicals.

Also solved by G. B. M. ZERR and J. F. SCHEFFER.

NOTE ON PROBLEM 391

BY LEWIS NEIKIRK, BOULDER, COLORADO.

The man starts at O moving in a *perfectly random* manner. After t seconds suppose him at P and that during the next instant dt he travels through ds to m at an angle θ with the line OP . Let $PM = dr = d\cos\theta = v\cos\theta dt$, since $ds = vdt$. He will escape from the desert if $\int dr > R$ (the radius) the limits of inte-